

On the conical refraction of hydromagnetic waves in plasma with anisotropic thermal pressure: general consideration

David Tsiklauri *

Physics Department, University of Cape Town, Rondebosch 7700, Republic of South Africa
(February 9, 2008)

A phenomenon analogous to the conical refraction well-known in the crystallooptics and crystalloacoustics is considered for the magnetohydrodynamical waves in a collisionless plasma with anisotropic thermal pressure. Imposing the most general condition for the existence of the phenomenon, angle of the conical refraction is calculated which appeared to be dependent on the ratio of the Alfvén velocity and sound speed measured in the perpendicular direction in respect to the external magnetic field. Feasible ways of experimental demonstration of the phenomenon are discussed and a novelty brought by the general consideration is outlined.

52.35.Bj, 42.15.Gs

I. INTRODUCTION

Recently, author of Ref.[1] showed, theoretically, that in the magnetized plasma with anisotropic thermal pressure [viz., a medium where linear magnetohydrodynamic (MHD) waves are described by well-known double adiabatic theory of Chew, Goldberger and Low [2]] there is a possibility of observation of the phenomenon of the conical refraction. The similar kind of phenomenon was observed long time ago (see, for example literature in Ref.[1]) for the light and acoustical waves in the biaxial crystals and therefore, it is very well established. For instance, in the biaxial crystals observation of the optical conical refraction is possible if the parallel, narrow beam of natural (non-polarized) light falls along the optical axis (binormal) of wave normals on a plate cut so that its two parallel faces are perpendicular to the binormal. While propagating through the crystal the beam will form an empty cone with permanently changing linear polarization. Thus, producing a circle on the photographic plate placed after the crystal.

In the Ref.[1] there were obtained the necessary conditions which ensure existence of the conical refraction in the medium under consideration. However, further investigation revealed that it is possible to impose much less restrictive conditions fulfillment of which ensures occurrence of the phenomenon. In the light of possible interest which plasma physics community may have to corroborate experimentally existence of the phenomenon, author find it worthwhile to outline here these novel conditions.

In the section II we derive the basic equations and outline new, general conditions for the existence of the phenomenon. While in the section III we pay more emphasis to the physics of the effect; discuss possible ways of its experimental corroboration along with its feasible applications for plasma diagnostics; and finally we address a novelty brought by the general consideration employed in the present paper.

II. MAIN CONSIDERATION

Let us consider the most general form of the dispersion relation for the MHD waves in magnetized plasma with anisotropic thermal pressure:

$$\omega^2 = \frac{k^2}{2\rho_0} \left[\frac{B_0^2}{4\pi} + p_{\perp}(1 + \delta \sin^2 \theta) + 2\alpha p_{\parallel} \cos^2 \theta \right. \\ \left. \pm \sqrt{\left(\frac{B_0^2}{4\pi} + p_{\perp}(1 + \delta \sin^2 \theta) - 2(\alpha + 1)p_{\parallel} \cos^2 \theta \right)^2 + 4p_{\perp}^2 \sin^2 \theta \cos^2 \theta} \right]. \quad (1)$$

*email: TSIKLAURI@physci.uct.ac.za

On leave from the Department of Theoretical Astrophysics, Abastumani Astrophysical Observatory, 2^a Kazbegi ave., Tbilisi 380060, Republic of Georgia.

The latter equation can be derived from the dispersion relation obtained in Ref.[3] while considering its nonrelativistic limit (i.e. neglecting displacement current and ultrarelativistic temperature effects), but retaining a freedom in choice of the polytropic indexes δ and α . Note, that in the derivation of Eq.(1) there were used following equations of state:

$$p_{\perp} = \text{const}_{\perp} \rho B^{\delta}, \quad (2)$$

$$p_{\parallel} = \text{const}_{\parallel} \rho \left(\frac{\rho}{B} \right)^{2\alpha}. \quad (3)$$

We gather from the Eqs.(2) and (3) that when $p_{\parallel} = p_{\perp} = p$ and $\alpha = \delta = 0$ the equations of state reduce to conventional one for isothermal, ideal gas $P \sim \rho$. For anisotropic pressure case, when $p_{\parallel} \neq p_{\perp}$ and $\alpha = \delta = 1$ they represent usual Chew, Goldberger and Low's equations of state [2]. Of course, taking into due account the two above limiting cases ($\alpha = \delta = 0$ and $\alpha = \delta = 1$) in Eq.(1) results in the correct expressions for the dispersion relations for the linear MHD waves in isotropic and anisotropic pressure plasmas accordingly, which can be found in any textbook of plasma physics.

For the sake of convenience we use further following notations [1]:

$$a = \frac{B_0}{\sqrt{4\pi\rho_0}}, \quad c_{\perp} = \sqrt{\frac{p_{\perp}}{\rho_0}}, \quad c_{\parallel} = \sqrt{\frac{p_{\parallel}}{\rho_0}}$$

where a stands for the Alfvén velocity, $c_{\perp}$ and $c_{\parallel}$ are the sound speeds measured in perpendicular and parallel directions in respect to the uniform, external magnetic field $\vec{B}_0$.

Now we can rewrite Eq.(1) in the following form:

$$V^2 = \frac{1}{2} \left[a^2 + c_{\perp}^2 (1 + \delta \sin^2 \theta) + 2\alpha c_{\parallel}^2 \cos^2 \theta \right. \\ \left. \pm \sqrt{\left(a^2 + c_{\perp}^2 (1 + \delta \sin^2 \theta) - 2(\alpha + 1)c_{\parallel}^2 \cos^2 \theta \right)^2 + 4c_{\perp}^4 \sin^2 \theta \cos^2 \theta} \right] \quad (4)$$

where $V = \omega/k$ denotes the phase velocity of the waves.

In Ref.[1] after this step, in order get desired result (theoretical demonstration of the existence of the conical refraction) author required fulfillment of the following relation between $c_{\parallel}$ and $c_{\perp}$: $c_{\perp} = \sqrt{2}c_{\parallel}$. Furthermore, according to Ref.[1] final necessary condition for occurrence of the effect was obtained as: $a = c_{\perp} = \sqrt{2}c_{\parallel}$. However, now let us restrict ourselves with much more weak condition:

$$a^2 + c_{\perp}^2 - 2(\alpha + 1)c_{\parallel}^2 = 0, \quad \text{or} \quad c_{\parallel}^2 = \frac{a^2 + c_{\perp}^2}{2(\alpha + 1)}. \quad (5)$$

Substituting in Eq.(4) the latter expression for $c_{\parallel}^2$ and introducing χ and ψ as

$$\chi^2 = a^2 + (1 + \delta)c_{\perp}^2 \quad \text{and} \quad \psi^2 = \frac{\alpha a^2 - [\delta(\alpha + 1) - \alpha]c_{\perp}^2}{\alpha + 1}$$

we can obtain:

$$V^2 = \frac{1}{2} \left[\chi^2 + \psi^2 \cos^2 \theta \pm \sqrt{\chi^4 + 2(2c_{\perp}^4 - \chi^4) \cos^2 \theta + (\chi^4 - 4c_{\perp}^4) \cos^4 \theta} \right] \quad (6)$$

Possessing the formula for the phase velocity of the wave it is easy to derive expression for the group velocity $\vec{u}$ via straightforward, but time consuming calculations:

$$\vec{u} = \frac{\partial \omega}{\partial \vec{k}} = \vec{N}V + k \frac{\partial V}{\partial \vec{k}} = \\ V \left[\frac{[\cos \theta / 2V^4] (\psi^2 T(\theta) + (2c_{\perp}^4 - \chi^4) + (\chi^4 - 4c_{\perp}^4) \cos^2 \theta)}{1 - C/V^4} \vec{A} + \right.$$

$$\left[\frac{[1/2V^4] \left((2V^2 - \psi^2 \cos^2 \theta) T(\theta) - (2c_\perp^4 - \chi^4) \cos^2 \theta - (\chi^4 - 4c_\perp^4) \cos^4 \theta \right)}{1 - C/V^4} \vec{N} \right] \quad (7)$$

where we introduced following notations: unit vectors $\vec{N} = \vec{k}/|\vec{k}|$ and $\vec{A} = \vec{B}_0/|\vec{B}_0|$; $T(\theta) \equiv 2V^2 - (\chi^2 + \psi^2 \cos^2 \theta)$ and

$$C \equiv (2\alpha + 1)[a^2 + c_\perp^2(1 + \delta \sin^2 \theta)] \left(\frac{a^2 + c_\perp^2}{2(\alpha + 1)} \right) \cos^2 \theta \\ - (2\alpha + 1) \left(\frac{a^2 + c_\perp^2}{2(\alpha + 1)} \right)^2 \cos^4 \theta - c_\perp^4 \sin^2 \theta \cos^2 \theta.$$

Here we must note that in Eq.(7) we used biquadratic equation

$$V^4 - (\chi^2 + \psi^2 \cos^2 \theta) V^2 + C = 0,$$

solution of which dispersion relation (6) actually is.

Let us describe a direction of the group velocity $\vec{u}$ by an angle φ between $\vec{u}$ and $\vec{B}_0$ (i.e., between $\vec{u}$ and $\vec{A}$). For the $\tan \varphi$ we may write then

$$\tan \varphi = \frac{\vec{u} \times \vec{A}}{\vec{u} \cdot \vec{A}} = \frac{(2V^2 - \psi^2 \cos^2 \theta) T(\theta) - (2c_\perp^4 - \chi^4) \cos^2 \theta - (\chi^4 - 4c_\perp^4) \cos^4 \theta}{(2V^2 + \psi^2 \sin^2 \theta) T(\theta) + (2c_\perp^4 - \chi^4) \sin^2 \theta + (\chi^4 - 4c_\perp^4) \cos^2 \theta \sin^2 \theta} \tan \theta. \quad (8)$$

Note, that $T(\theta) \rightarrow 0$ when $\theta \rightarrow 0$.

The latter equation determines the direction of the group velocity $\vec{u}$ whereas its modulus can be calculated by expression:

$$|\vec{u}| = V / \cos(\varphi - \theta) \quad (9)$$

It is worth of remarking that the Eq.(9) is a consequence of the following expression $\vec{u} \cdot \vec{N} = V$ which can be readily shown by multiplying Eq.(7) scalarly by $\vec{N}$.

Now, it can be easily shown that the expressions for group velocity and $\tan \varphi$ [Eqs. (7) and (8) correspondingly] exhibit uncertain behaviour as $0/0$ when $\theta \rightarrow 0$. Removing of the uncertainty (for the sake of brevity we omit, here, several steps of this task) results in:

$$\tan \varphi(+0) = \lim_{\theta \rightarrow 0} \tan \varphi(\theta) = \pm \left(\frac{\alpha + 1}{2\alpha + 1} \right) \frac{1}{1 + (a/c_\perp)^2} \equiv \pm \left(\frac{\alpha + 1}{2\alpha + 1} \right) \Gamma^2, \quad (10)$$

$$|\vec{u}(+0)| = \sqrt{\left[1 + \left(\frac{\alpha + 1}{2\alpha + 1} \right)^2 \Gamma^4 \right] \left(\frac{2\alpha + 1}{\alpha + 1} \right) \sqrt{\frac{a^2 + c_\perp^2}{2}}}. \quad (11)$$

In Eq.(10) sign "+" and "-" correspond to the slow and fast MHD waves. It is worthwhile to note that Eqs.(10) and (11) are, of course, in agreement with previous results [1,4]. If we consider the isotropic pressure limit ($p_\parallel = p_\perp = p$, $c_\parallel = c_\perp = c$, and $\alpha = \delta = 0$) and require $a^2 = c^2$ [4] then Eqs.(10) and (11) reduce to the form:

$$\tan \varphi(+0) = \pm \frac{1}{2} \quad \text{and} \quad |\vec{u}(+0)| = \frac{\sqrt{5}}{2} c = \frac{\sqrt{5}}{2} a \quad (12)$$

as it was actually obtained in Ref.[4]. Whereas, applying the condition imposed in Ref.[1] $a = c_\perp = \sqrt{2} c_\parallel$ we arrive at

$$\tan \varphi(+0) = \pm \frac{1}{3} \quad \text{and} \quad |\vec{u}(+0)| = \sqrt{\frac{5}{3}} c_\perp = \sqrt{\frac{5}{3}} a. \quad (13)$$

We must admit, here, that in the Eq.(9) for the modulus of the group velocity from the Ref.[1] factor of $\sqrt{\frac{3}{2}}$ is missing. Apparently, substituting $V(\theta = 0) = \sqrt{\frac{3}{2}} \cdot a = \sqrt{\frac{3}{2}} \cdot c_{\perp}$ and $\cos \varphi = \sqrt{10}/3$ in the Eq.(9) (of the present paper) results in Eq.(13). This should not cause a confusion, as it does not affect the main result of the Ref.[1] — theoretical demonstration of possibility of the occurrence of the conical refraction in the anisotropic plasma plus calculation of the angle of the conical refraction, which is, actually, an experimentally observable physical quantity. We see that the results obtained in this paper are consistent with previous contributions [1,4]. However, it must be emphasized that more general results given by Eqs.(10) and (11) bring more diversion in the predicted phenomenon (see below). Besides, from technical point of view it should be much easier to produce the plasma satisfying condition (5) obtained in this paper, rather than condition $a = c_{\perp} = \sqrt{2}c_{\parallel}$ requested in Ref.[1].

III. DISCUSSION

The above consideration shows that when the wave normal $\vec{N}$ tends to reach a direction parallel to $\vec{B}_0$ (i. e. when θ tends to zero), vector of group velocity $\vec{u}$ does not take the same direction, but its limiting position (when $\theta \rightarrow 0$) is the surface of an empty cone with opening angle $2\varphi(+0) \equiv 2 \arctan[\varphi(+0)]$ given by Eq.(10). When θ is zero exactly (which is quite unlikely from the experimental point of view as all waves produced in a laboratory may have only finite angle of aperture) then $\vec{u}$ is strictly parallel to $\vec{B}_0$. The discontinuous behavior exhibited by the vector $\vec{u}$ can be explained by means of existence of so called [1,4] angular point which in the case of the anisotropic plasma revealed to exist when $a^2 + c_{\perp}^2 - 2(\alpha + 1)c_{\parallel}^2 = 0$. Thus, when $a^2 + c_{\perp}^2 - 2(\alpha + 1)c_{\parallel}^2 = 0$ the angular point on the surfaces formed by the phase velocities (both fast and slow MHD waves) is located in the point of intersection of the ones with the direction of the magnetic field $\vec{B}_0$. Which, in turn leads to the discontinuous behaviour of $\vec{u}$ and causes the effect. In other words, as it was mentioned in Ref.[1], the angular point exists when phase velocities of the slow and fast MHD waves do coincide. We observe from Eq. (6) [again, which was obtained from general dispersion relation (4) imposing condition (5)] that when $\theta \rightarrow 0$ slow ("−") and fast ("+") branches of phase velocity equal to each other (viz., the square root vanishes), which results in an uncertain behavior of its derivative by $\vec{k}$ i.e. group velocity.

It is interesting to note that in differ to the result of Ref.[1], now imposing less restrictive condition (5) the angle of the conical refraction is not a *fixed* number but it depends [via Eq.(10)] on the ratio $a^2/c_{\perp}^2$. Which, in turn, depends on the plasma medium physical parameters such as density and the magnitude of the exerted external magnetic field $\vec{B}_0$.

As regards possible experimental corroboration of the effect, we may state the following: consider two collisionless plasma media contained in a uniform external magnetic field $\vec{B}_0$. The media are separated by means of the insulating plane perpendicular to the $\vec{B}_0$. Let us choose physical parameters in the second medium as to satisfy the condition (5). If we place a diaphragm between the first and second media then a narrow beam of MHD waves propagating from the first medium towards the second one, just after passing the diaphragm will deflect from the direction of $\vec{B}_0$ and form an empty cone with angle of opening $2\varphi(+0)$.

It is worthwhile to note that one of the possible applications of the effect of the conical refraction may be its usage in plasma diagnostics. The condition (5) explicitly depends on the magnitude of external magnetic field $\vec{B}_0$. So, if in the above proposed experimental setup, in the second medium the condition (5) is not satisfied a detector of MHD waves placed just opposite to the diaphragm (see Fig.1) will fairly detect the waves. However, with variation of the magnitude of the exerted magnetic field we may achieve fulfillment of the condition (5) which will result in a steep drop of the detector's indication, as due to the effect of the conical refraction the MHD waves will deflect from the $\theta = 0$ axis and form an empty cone with opening angle $2\varphi(+0)$, thus avoiding arrival of the waves into the detector (see Fig. 2). Since the condition (5) explicitly depends on the external magnetic field, in the event of occurrence of the phenomenon it must be possible to measure its magnitude if the thermodynamical parameters (pressure and density) of the plasma are known or vice versa. Therefore, we conclude that presumably this effect can be used in plasma diagnostics.

Besides, of a purely fundamental interest which experimental demonstration of the phenomenon might have, probably, this effect will find its applications in practice, since through variation of external parameters (viz., varying the ratio $a^2/c_{\perp}^2$) it is possible to regulate opening angle of the empty conical beam of the MHD waves. Which means, that, for instance, with simple from technical point of view variation of the magnitude of the external magnetic field it is possible to regulate a direction of propagation of the energy carried by the MHD waves. Because, it is known that the energy propagates along the direction of the wave's group velocity vector. Therefore, one should not exclude possible applications of this novel effect (revealed by the general consideration employed in this paper) in the plasma devices like MHD generators are.

Finally, we would like to emphasize on principal difference between the effect predicted in this paper and the effect of the conical refraction occurring in the optically and acoustically biaxial crystals. First, the most evident difference is that the phenomenon can take place in media with qualitatively different physical properties (magnetized plasma with isotropic [4] or anisotropic [1] pressure on the one hand and the biaxial crystals on the other). Second, in the case of biaxial crystals, let us take for concreteness optical conical refraction, mathematical basis for theoretical formulation of the effect constitutes Maxwell's equations along with material equations and in addition *electrical* anisotropy of the medium is assumed, i.e. such substances (crystals) are considered whose electrical excitations depend on the direction of the electric field [5]. Whereas in our case anisotropy in the medium is brought by the external *magnetic* field and Euler's equation for the magnetized, perfectly conducting plasma is used. The only similarity of these two phenomena is that the equations which relate the phase velocity of the waves with their direction of propagation in respect to the anisotropy axis (optical axis in one case and the external magnetic field in the other) — Fresnel's equation of wave normals [5] and our dispersion relation (4) (including its special subcases [1] and [4]) have the same mathematical structure. Thus, geometrical shape of the surface formed by the group velocity vectors when the phenomenon takes place is qualitatively the same — surface of the empty cone.

IV. ACKNOWLEDGEMENTS

This research was supported in part by the Research Scholarship of the University of Cape Town and by South African Foundation for Research Development (FRD).

-
- [1] D. Tsiklauri, Phys. Plasmas, **3**, 800, (1996).
 - [2] G. Chew, M. Goldberger, and F. Low, Proc. R. Soc. London ser. **A** **236**, 112, (1956).
 - [3] E. G. Tsikarishvili and A. D. Rogava, Phys. Rev. E **50**, 3050, (1994).
 - [4] D.V. Sivukhin, Magnitnaya Gidrodinamica, **1**, 35, (1966). (in Russian)
 - [5] M. Born and E. Wolf, *Principles of Optics*, (Pergamon Press, 1980).




FIG. 1. Sketch of experimental set up, which may be used for demonstration of the effect of the conical refraction (see details in the text). The line with arrows represents the narrow, parallel beam of MHD waves. Note, that when in the medium II the condition $a^2 + c_{\perp}^2 - 2(\alpha + 1)c_{\parallel}^2 = 0$ is not satisfied the beam goes into detector without experiencing any changes while propagating through the second medium.




FIG. 2. The same as in Fig.1. Note, that when in the medium II the condition $a^2 + c_{\perp}^2 - 2(\alpha + 1)c_{\parallel}^2 = 0$ is fulfilled, the beam due to the effect of the conical refraction deflects from its initial direction ($\theta = 0$) and forms an empty cone with opening angle $2\varphi(+0) \equiv 2 \arctan[\varphi(+0)]$, thus avoiding arrival of the MHD waves into detector's receiver.